

## Ammonia Induces Settlement Behavior in Oyster Larvae

S. L. COON<sup>1</sup>, M. WALCH<sup>2,3</sup>, W. K. FITT<sup>4</sup>, R. M. WEINER<sup>2,3</sup>, AND D. B. BONAR<sup>1,3</sup>

<sup>1</sup>*Department of Zoology, <sup>2</sup>Department of Microbiology, and <sup>3</sup>Center of Marine Biotechnology, University of Maryland, College Park, Maryland 20742, and <sup>4</sup>Department of Zoology, University of Georgia, Athens, Georgia 30602*

**Abstract.** Oyster larvae exposed to solutions of  $\text{NH}_4\text{Cl}$  exhibit stereotypical settlement behavior similar to that which normally precedes cementation and metamorphosis. Un-ionized ammonia is the active chemical species. At  $\text{pH} = 8.0$ , the threshold concentration of  $\text{NH}_4\text{Cl}$  ( $\text{pH} = 8.0$ ) for newly competent larvae is  $2.5 \text{ mM}$ ; maximum activity is at  $7.9 \text{ mM}$ , corresponding to calculated  $\text{NH}_3$  concentrations of  $100 \mu\text{M}$  and  $310 \mu\text{M}$ , respectively. Induction of settlement behavior is rapid, with  $>90\%$  of larvae exposed to  $310 \mu\text{M}$   $\text{NH}_3$  responding within less than 5 min. After 15 to 30 min, larvae become habituated to  $\text{NH}_3$  and resume swimming so that the percent exhibiting settlement behavior after 30 min is  $<10\%$ . Other weak bases, such as methylamine and trimethylamine, induce similar behavior suggesting that  $\text{NH}_3$  acts by increasing intracellular  $\text{pH}$ . Evidence that  $\text{NH}_3$  and L-3,4-dihydroxyphenylalanine (L-DOPA) induce settlement behavior through different mechanisms is presented. Ammonia may be a natural environmental cue that promotes oyster settlement behavior and, ultimately, recruitment.

### Introduction

Many marine invertebrates, including oysters, have planktonic larvae that are recruited preferentially to habitats suitable for subsequent survival (Thorson, 1950). Recruitment of invertebrate larvae often involves a stereotyped series of search and crawl behaviors that is called settlement, followed by a morphogenetic phase called

metamorphosis (Burke, 1983). The settlement behavior of oyster larvae has been well characterized and includes swimming with the foot extended forward followed by a series of increasingly localized crawling maneuvers (Prytherch, 1932; Cranfield, 1973; Coon *et al.*, 1985). If the habitat in which the larva has settled is suitable, the larva will cement permanently to the substratum and metamorphose. Settlement is reversible and does not necessarily culminate in metamorphosis once initiated; if the habitat is unsuitable, the larva may resume swimming and repeat the process elsewhere.

Invertebrate larvae are often induced to settle and metamorphose by environmental cues, typically chemical, associated with the adult habitat (Crisp, 1974; Chia and Rice, 1978). Microbial films play an important role in the development of many invertebrate assemblages (Zobell and Allen, 1935; Meadows and Campbell, 1972; Scheltema, 1974; Bonar *et al.*, 1986). Both soluble and surface-associated bacterial products are important in recruiting invertebrate larvae to surfaces containing bacterial films (Wilson, 1955; Scheltema, 1961; Gray, 1967; Muller, 1973; Neumann, 1979; Kirchman *et al.*, 1982), although some larvae prefer unfiled surfaces (Crisp and Ryland, 1960). A bacterium, *Alteromonas colwelliana* (originally called LST), was found to enhance the recruitment of oyster larvae to colonized substrates (Weiner *et al.*, 1985; 1989). Supernatants from cultures of *A. colwelliana*, as well as other bacteria, contain one or more soluble factors that induce settlement behavior in oyster larvae of the genus *Crassostrea*. Preliminary studies showed that the soluble inducer has a low molecular weight ( $<300$  daltons), and that supernatants have increased inductive potency commensurate with the age of the bacterial culture (Fitt *et al.*, 1990).

Received 4 June 1990; accepted 25 September 1990.

Contribution #141 from the Center of Biotechnology, Marine Biotechnology Institute, University of Maryland.

Experiments reported in this paper demonstrate that solutions of  $\text{NH}_4\text{Cl}$  induce settlement behavior, and that  $\text{NH}_3$ , not  $\text{NH}_4^+$ , is the active chemical species. Additional experiments further explore the relationship between the mechanism of  $\text{NH}_3$  induction and induction of settlement behavior by L-3,4-dihydroxyphenylalanine (L-DOPA), another known soluble inducer of oyster settlement behavior (Coon *et al.*, 1985, 1990). Preliminary results of this work have been presented (Coon *et al.*, 1988; Bonar *et al.*, 1990).

## Materials and Methods

### *Obtaining and maintaining larvae*

Larvae of the Pacific oyster, *Crassostrea gigas*, were obtained from the Coast Oyster Company of Quilcene, Washington, and maintained in the laboratory (Coon *et al.*, 1990). Larvae were used within one week of arrival.

### *Bioassay procedure*

Experiments were conducted as previously described (Coon *et al.*, 1990). Aliquots of 20–50 larvae were assayed in 24-well tissue culture plates (Falcon #3047) in a final volume of 1.0 ml. Antibiotics were not used, but all experiments were conducted in 0.2  $\mu\text{m}$  filtered seawater. Each treatment was duplicated or triplicated, and results are expressed as the mean  $\pm$  standard error. All chemicals were obtained from Sigma Chemical Company (St. Louis, Missouri).

Larval settlement behavior was defined as in Coon *et al.* (1990), the basic criterion being active foot extension beyond the ventral margin of the shell. Behavior in each well was monitored with a dissecting microscope for 30 s at the times noted. The length of each experiment was between 30 and 40 min, as noted.

Statistical tests were performed on arcsine-transformed data using a one-way analysis of variance (ANOVA) within time points. The ANOVA was followed by a Student-Newman-Keuls pair-wise comparisons test when significant differences were detected (Zar, 1974). Differences were considered significant if  $P < 0.05$ .

### *Effects of ammonia, ammonium, and pH*

Larvae were exposed to a range of concentrations of  $\text{NH}_4\text{Cl}$ , in the first series of experiments. Stock solutions of  $\text{NH}_4\text{Cl}$  were made in seawater at twice the final concentration and adjusted to  $\text{pH} = 8.0$  with  $\text{NaOH}$ . At the beginning of each bioassay, 0.5 ml of stock solution was added to an equal volume of seawater ( $\text{pH} = 8.0$ ) containing swimming larvae. This experiment was repeated using  $(\text{NH}_4)_2\text{SO}_4$  and other chloride salts ( $\text{NaCl}$ ,  $\text{KCl}$ ) at concentrations up to 10 mM.


Approximately 96% of the total ( $\text{NH}_3 + \text{NH}_4^+$ ) in seawater at  $\text{pH} = 8.0$  is present as the ammonium ion,  $\text{NH}_4^+$  (Bower and Bidwell, 1978). To determine whether  $\text{NH}_3$  or  $\text{NH}_4^+$  was the active chemical species, larval settlement responses were observed while the concentrations of  $\text{NH}_3$  and  $\text{NH}_4^+$  were varied under two different regimes. In the first,  $\text{pH}$  was held constant at 8.0 and the total ( $\text{NH}_3 + \text{NH}_4^+$ ), as  $\text{NH}_4\text{Cl}$ , was varied as described for the initial experiments above. In the second regime, total ( $\text{NH}_3 + \text{NH}_4^+$ ) was held constant at 5.0 mM and the proportion of  $\text{NH}_3$  to  $\text{NH}_4^+$  was varied by altering the  $\text{pH}$ . The absolute concentrations of  $\text{NH}_3$  and  $\text{NH}_4^+$  were calculated by means of a hydrolysis constant for ammonium ion in seawater of  $\text{pK}_a^s = 9.39$ , at 30‰ salinity, 23°C and 1 atm pressure (Bower and Bidwell, 1978).

Because ammonia is a weak base ( $\text{pK}_a = 9.25$ ), its effects might result from an increase in intracellular  $\text{pH}$  ( $\text{pH}_i$ ). Therefore, two other weak bases, methylamine ( $\text{pK}_a = 10.7$ ) and trimethylamine ( $\text{pK}_a = 9.81$ ), were tested for their ability to induce settlement behavior. The inductive activities of these two compounds, along with those of  $\text{NH}_4\text{Cl}$ , were investigated according to the original protocol described above; concentration was varied while the  $\text{pH}$  remained constant at 8.0.

### *Relationship of $\text{NH}_3$ -induction to L-DOPA-induction of settlement behavior*

To determine whether  $\text{NH}_3$  and L-DOPA induce settlement behavior through the same mechanisms, we tested sulpiride, a dopaminergic receptor antagonist (Stoof and Keabian, 1984) and potent inhibitor of L-DOPA-induced settlement behavior (Coon and Bonar, 1987), for its ability to block  $\text{NH}_3$ -induced settlement behavior. Ammonium chloride stock solutions were made 10 times the final concentration in filtered seawater and adjusted to  $\text{pH} = 8.0$ . Solutions of L-DOPA and sulpiride were made 10 times their final concentrations in 0.002 *N* HCl. All larvae were pre-incubated in either 100  $\mu\text{M}$  sulpiride or seawater for 12 min, then exposed to either 10 mM  $\text{NH}_4\text{Cl}$  or 100  $\mu\text{M}$  L-DOPA. The effects of seawater and 0.002 *N* HCl were appropriately controlled. The  $\text{pH}$  of the final solutions was 7.7, yielding a calculated  $\text{NH}_3$  concentration of 270  $\mu\text{M}$ .

In other experiments, larvae that were "habituated" to  $\text{NH}_3$  (see Results) were tested to see whether they would still respond to L-DOPA. Larvae were exposed to 5.0 mM  $\text{NH}_4\text{Cl}$  for 18 min until they began to habituate. They were then removed, rinsed, and exposed to either: (1) 100  $\mu\text{M}$  L-DOPA; (2) fresh 5.0 mM  $\text{NH}_4\text{Cl}$ ; (3) the  $\text{NH}_4\text{Cl}$  solution from which they had just been removed; or (4) filtered seawater. Control groups were pre-exposed to filtered seawater instead of  $\text{NH}_4\text{Cl}$ , then rinsed and put in



**Figure 1.** Percentages of *Crassostrea gigas* larvae exhibiting settlement behavior as a function of length of time exposed to specified concentrations of  $\text{NH}_4\text{Cl}$  at  $\text{pH} = 8.0$ . Calculated  $\text{NH}_3$  concentrations are shown in brackets for reference. Data are means of duplicates followed through time. Controls contained only filtered seawater ( $\text{pH} = 8.0$ ).

the same four treatments. In addition, some larvae were left in the  $\text{NH}_4\text{Cl}$  solution without rinsing, and were exposed to either: (1) the addition of another 5.0 mM  $\text{NH}_4\text{Cl}$ ; (2) the addition of 100  $\mu\text{M}$  L-DOPA; or (3) no additional treatment. Larval settlement behavior was monitored for an additional 39 min.

## Results

### *Ammonium chloride induces settlement behavior*

Ammonium chloride induced high levels of settlement behavior in oyster larvae (Fig. 1). The percentage of larvae exhibiting settlement behavior reached >90% within 5 min of exposure to an  $\text{NH}_4\text{Cl}$  solution of 7.9 mM at  $\text{pH} = 8.0$ . Responses to higher concentrations of  $\text{NH}_4\text{Cl}$  are not shown because larvae in these solutions exhibited reduced activity levels after short exposures. Between 2.5 and 7.9 mM ( $\text{pH} = 8.0$ ), the larval response to  $\text{NH}_4\text{Cl}$  was concentration dependent. As the  $\text{NH}_4\text{Cl}$  concentration increased, the percentage of larvae exhibiting settlement behavior increased, and the length of time required for the maximum percentage of larvae to respond decreased. Following the maximum larval response, the percentage of larvae continuing to exhibit settlement behavior rapidly declined so that 30 min after the initial exposure, almost all the larvae had "habituated" to the  $\text{NH}_4\text{Cl}$  solutions and had resumed normal swimming. No subsequent metamorphosis was observed after 24 to 48 h.

Larvae also exhibited high levels of settlement behavior in response to  $(\text{NH}_4)_2\text{SO}_4$ , indicating that either  $\text{NH}_3$  or  $\text{NH}_4^+$  was the active chemical species. This was corrob-

orated by the observation that larval settlement behavior was not induced by  $\text{Cl}^-$ —as NaCl or KCl—at concentrations comparable to inductive  $\text{NH}_4\text{Cl}$  solutions (data not shown). Methylamine and trimethylamine, which are weak bases like  $\text{NH}_3$ , induced high levels of oyster settlement behavior (Table I).

### *The active species is $\text{NH}_3$ rather than $\text{NH}_4^+$*

Ammonia has a  $\text{pK}_a^s$  of 9.39 in seawater and its calculated speciation as a function of  $\text{pH}$  is shown in Figure 2A. As the  $\text{pH}$  of the  $\text{NH}_4\text{Cl}$  solution decreases from 8.0 to 7.0, which is within the physiological tolerance range for oyster larvae, the  $\text{NH}_3$  concentration changes much more dramatically (89.6% decrease) than the  $\text{NH}_4^+$  concentration (3.6% increase) (Fig. 2B). Theoretically, the chemical species,  $\text{NH}_3$  or  $\text{NH}_4^+$ , to which the larvae are responding, would have the same dose-response curve, whether the concentrations are adjusted by varying the  $\text{NH}_4\text{Cl}$  concentration under constant  $\text{pH}$ , or by keeping the  $\text{NH}_4\text{Cl}$  concentration constant and varying the  $\text{pH}$ .

This experiment shows that the maximal percentage of larvae exhibiting settlement behavior in response to  $\text{NH}_3$  was independent of the regime used to vary the  $\text{NH}_3$  concentration (Fig. 3A). The difference between these two curves represents less than 0.1  $\text{pH}$  unit, which was within experimental error. In contrast, the larval response to  $\text{NH}_4^+$  was highly dependent on the regime used to vary the  $\text{NH}_4^+$  concentration; the larval response increased with increasing  $\text{NH}_4^+$  concentration when the  $\text{pH}$  was held constant while the  $\text{NH}_4\text{Cl}$  concentration was varied, but the larval response decreased with increasing  $\text{NH}_4^+$  concentration when the  $\text{NH}_4\text{Cl}$  was held constant while the  $\text{pH}$  was varied (Fig. 3B). These results indicate that, in these solutions,  $\text{NH}_3$ , not  $\text{NH}_4^+$ , was the active chemical species inducing settlement behavior in oyster larvae.

### *$\text{NH}_3$ and L-DOPA induce settlement behavior through different mechanisms*


The dopaminergic antagonist, sulpiride, blocked the ability of L-DOPA to induce settlement behavior (Fig.

**Table I**

*Maximal percentage of oyster larvae exhibiting settlement behavior in response to exposure to weak bases at  $\text{pH} = 8.0$*

	$\text{pK}_a$	1.0 mM	3.3 mM	10 mM
$\text{NH}_4\text{Cl}$	9.25	3.2 $\pm$ 0.4	51.2 $\pm$ 4.0	90.3 $\pm$ 2.3
Methylamine	10.7	4.4 $\pm$ 1.4	47.6 $\pm$ 14.2	94.6 $\pm$ 2.4
Trimethylamine	9.81	0	19.7 $\pm$ 7.6	91.6 $\pm$ 5.4

Data are means of duplicates  $\pm$  standard error.



**Figure 2.** Calculated effect of pH on NH<sub>3</sub> and NH<sub>4</sub><sup>+</sup> speciation. (A) Percentage contribution to the total (NH<sub>3</sub> + NH<sub>4</sub><sup>+</sup>) by each species as a function of pH.  $pK_a^s = 9.39$ . (B) Percentage change in NH<sub>3</sub> and NH<sub>4</sub><sup>+</sup> as the pH of the solution drops from pH = 8.0 to the specified value. Actual calculated changes in speciation are tabulated for clarity.


4A) but did not block the ability of NH<sub>3</sub> to induce settlement behavior (Fig. 4B). However, two small effects of sulphuride on NH<sub>3</sub>-induced settlement behavior were noted: (1) settlement behavior was more rapidly expressed; and (2) the maximum percentage of larvae induced to exhibit settlement behavior was slightly lower (*t*-test;  $P < 0.1$ ). The differential effects of sulphuride on the abilities of NH<sub>3</sub> and L-DOPA to induce settlement behavior indicate that NH<sub>3</sub> functions through a mechanism that does not require the dopaminergic receptor involved in the induction of settlement behavior by L-DOPA (Coon and Bonar, 1987).

The effects of NH<sub>3</sub> and L-DOPA are not completely independent. Although larvae that had habituated to NH<sub>3</sub> were almost completely refractory to fresh NH<sub>3</sub>, they could still respond to L-DOPA (Fig. 5). However, fewer of these larvae exhibited settlement behavior, and they responded more slowly to L-DOPA than larvae that had not been habituated to NH<sub>3</sub>. The larvae also showed an attenuated response to L-DOPA in additional treatments during which they were left in the presence of NH<sub>3</sub> when


L-DOPA was added (data not shown). The small, transient, increase in settlement behavior following transfer to a new NH<sub>4</sub>Cl solution was an artifact of the procedure.

## Discussion

This study demonstrates that NH<sub>3</sub> in the surrounding medium induces oyster larvae to exhibit settlement behavior. The onset of settlement behavior is rapid, high percentages of larvae are induced to behave, and the larvae quickly resume swimming without cementing to the plastic culture plates (a suboptimal settlement surface) in which the experiments were conducted. The results also indicate that, although NH<sub>3</sub> and L-DOPA induce similar settlement behaviors, the biochemical mechanisms by which they do so are different.



**Figure 3.** Maximal percentages of *Crassostrea gigas* larvae exhibiting settlement behavior as a function of the concentration of either NH<sub>3</sub> or NH<sub>4</sub><sup>+</sup>. In one regime, pH was held constant while the NH<sub>4</sub>Cl concentration was varied (data calculated from Fig. 1); in the other regime, NH<sub>4</sub>Cl concentration was held constant while the pH was varied. The concentrations of NH<sub>3</sub> and NH<sub>4</sub><sup>+</sup> were calculated from pH values and NH<sub>4</sub>Cl concentration. (A) Larval response as a function of the calculated concentration of NH<sub>3</sub> under the two regimes. (B) Larval response as a function of NH<sub>4</sub><sup>+</sup> under the two regimes. Data are means of duplicates.




**Figure 4.** Percentages of *Crassostrea gigas* larvae exhibiting settlement behavior as a function of length of the duration of their exposure to  $\text{NH}_3$  or L-DOPA in the presence of sulpiride. Sulpiride is a dopaminergic receptor antagonist. (A) Effect of sulpiride on the ability of L-DOPA to induce settlement behavior. (B) Effect of sulpiride on the ability of  $\text{NH}_3$  to induce settlement behavior. Larvae were pre-incubated in sulpiride ( $100 \mu\text{M}$ ) or seawater for 12 min, then L-DOPA ( $100 \mu\text{M}$ ),  $\text{NH}_4\text{Cl}$  ( $10 \text{ mM}$ ), sulpiride ( $100 \mu\text{M}$ ) or seawater were added as indicated by the arrows. Data are means  $\pm$  standard error of triplicates followed through time. For each time point, treatments with the same letter, or no letter are not significantly different from each other. Only statistics for relevant time points are shown for clarity.

Ammonia, as a by-product of protein catabolism, is excreted by most marine bacteria and animals (Campbell, 1970; Billen, 1984). Therefore, in areas of high biological activity and reduced mixing (such as in boundary layers near surfaces),  $\text{NH}_3$  might reach levels high enough to induce settlement behavior in oyster larvae. Total ( $\text{NH}_3 + \text{NH}_4^+$ ) concentrations of  $10 \text{ mM}$  have been reported in interstitial waters from marine sediments (Bruland, 1983). Stevens (1983) found that total ( $\text{NH}_3 + \text{NH}_4^+$ ) concentrations in association with oyster reefs may reach greater than  $200 \mu\text{M}$  in sediment waters and  $3 \mu\text{M}$  in overlying waters 10 cm above the sediment interface. The presence of high levels of  $\text{NH}_3$  in the environment, the rapid induction of settlement behavior by  $\text{NH}_3$ , and the

quick reversibility of inductive effects of  $\text{NH}_3$ , strongly suggest that  $\text{NH}_3$  is a natural environmental cue for recruitment of oyster larvae. Its actual involvement in larval recruitment, however, has not yet been demonstrated.

If  $\text{NH}_3$  is a natural inducer of settlement behavior, then it must be a relatively non-specific indicator of biologically rich environments. Ammonia alone could not account for the specificity observed in natural oyster settlement and metamorphosis. We hypothesize that  $\text{NH}_3$  acts as a chemokinetic agent that induces settlement behavior once a threshold concentration is encountered, bringing oyster larvae into contact with substrates and other potential contact-dependent and soluble cues (*c.f.* Crisp, 1974). Once settlement behavior has been initiated, oyster larvae rely on other cues from the environment to indicate that the habitat is suitable for cementation and metamorphosis. If these secondary cues are not present, the larvae habituate to  $\text{NH}_3$  and swim away. This scenario is consistent with models and observations of oyster settlement (Prytherch, 1934; Cranfield, 1973; Coon *et al.*, 1985; Weiner *et al.*, 1989; Coon *et al.*, 1990).

Although the mechanism by which  $\text{NH}_3$  induces settlement behavior in oysters is unknown,  $\text{NH}_3$  acts by increasing pH, in other invertebrate systems (Boron and DeWeer, 1976; Roos and Boron, 1981; Dube and Guerrier, 1982; Ward *et al.*, 1983; Bibring *et al.*, 1984; Williams *et al.*, 1984; Busa, 1986). Weak bases, such as  $\text{NH}_3$ , raise pH, by penetrating the cell membrane as the uncharged



**Figure 5.** Percentages of *Crassostrea gigas* larvae exhibiting settlement behavior as a function of length of time exposed to various regimes of  $\text{NH}_4\text{Cl}$  and L-DOPA. Larvae were exposed to either  $\text{NH}_4\text{Cl}$  ( $5 \text{ mM}$ ) or seawater for 18 min, then removed (downward pointing arrow) and put into either  $\text{NH}_4\text{Cl}$  ( $5 \text{ mM}$ ) or L-DOPA ( $100 \mu\text{M}$ ). Data are means  $\pm$  standard error of triplicates followed through time. For each time point, treatments with the same letter, or no letter are not significantly different from each other. Only statistics for relevant time points are shown for clarity.

species, then reprotonating in the cytoplasm (Roos and Boron, 1981). The induction of settlement behavior by other weak bases, such as methylamine and trimethylamine, is consistent with  $\text{NH}_3$  acting by increased intracellular alkalization. An increase in  $\text{pH}_i$  would not be expected to be cell-type specific and so may affect a diverse range of cell types in the larvae. Larvae of the hydroid, *Hydractinia*, are induced to metamorphose by  $\text{NH}_4^+$ , not  $\text{NH}_3$ , through a mechanism that may involve regulation of intracellular transmethylation rather than  $\text{pH}_i$  (Berking, 1988).

Whatever its mode of action, induction of settlement behavior by  $\text{NH}_3$  clearly involves a mechanism different from that of L-DOPA induction, though they are probably related. Ammonia and L-DOPA have different time courses. Larvae respond quickly to  $\text{NH}_3$ , then soon habituate to it; in contrast, larvae respond more slowly to L-DOPA, and the effects are longer lasting. Induction of settlement behavior by  $\text{NH}_3$  is not mediated through the same dopaminergic receptors required for induction by L-DOPA, but is effected slightly by blocking these receptors with sulphiride. Conversely, larvae habituated to  $\text{NH}_3$  can still respond to L-DOPA but to a lesser degree. There may be some interaction between  $\text{pH}_i$  and signal transduction through the dopaminergic receptors. Further experiments are underway to resolve the mechanism of  $\text{NH}_3$ -induction.

### Acknowledgments

The authors gratefully acknowledge the Coast Oyster Company of Quilcene, Washington, for providing the oyster larvae for this study. This research was supported by the National Science Foundation (PCM 831678), Maryland Department of Natural Resources (S-124-88-008), Maryland Sea Grant (NOAA-NA-86AAD-SG006) and the Maryland Industrial Partnership Program (#119.23).

### Literature Cited

- Berking, S. 1988. Ammonia, tetramethylammonium, barium and amiloride induce metamorphosis in the marine hydroid *Hydractinia*. *Roux's Arch. Dev. Biol.* **197**: 1-9.
- Bibring, T., J. Baxandall, and C. C. Harter. 1984. Sodium-dependent pH regulation in active sea urchin sperm. *Dev. Biol.* **101**: 425-435.
- Billen, G. 1984. Heterotrophic utilization and regeneration of nitrogen. Pp 313-355 in *Heterotrophic Activity in the Sea*, J. E. Hobbie, and P. J. leB. Williams, eds., Plenum Press, New York.
- Bonar, D. B., S. L. Coon, M. Walch, R. M. Weiner, and W. Fitt. 1990. Control of oyster settlement and metamorphosis by endogenous and exogenous chemical cues. *Bull. Mar. Sci.* **46**: 484-498.
- Bonar, D. B., R. M. Weiner, and R. R. Colwell. 1986. Microbial-invertebrate interactions and potential for biotechnology. *Microb. Ecol.* **12**: 101-110.
- Boron, W. F., and P. DeWeer. 1976. Intracellular pH transients in squid giant axons caused by  $\text{CO}_2$ ,  $\text{NH}_3$ , and metabolic inhibitors. *J. Gen. Physiol.* **67**: 91-112.
- Bower, C. E., and J. P. Bidwell. 1978. Ionization of ammonia in seawater: effects of temperature, pH, and salinity. *J. Fish. Res. Board Can.* **35**: 1012-1016.
- Bruland, K. W. 1983. Trace elements in seawater. Pp. 157-220 in *Chemical Oceanography*, J. P. Riley, and G. Skirrow, eds., Academic Press, New York.
- Burke, R. D. 1983. The induction of marine invertebrate larvae: stimulus and response. *Can. J. Zool.* **61**: 1701-1719.
- Busa, W. B. 1986. Mechanisms and consequences of pH-mediated cell regulation. *Ann. Rev. Physiol.* **48**: 389-402.
- Campbell, J. W., ed. 1970. *Comparative Biochemistry of Nitrogen Metabolism 1. The Invertebrates*. Academic Press, New York. 493 pp.
- Chia, F.-S., and M. E. Rice, eds. 1978. *Settlement and Metamorphosis of Marine Invertebrate Larvae*. Elsevier, New York. 290 pp.
- Coon, S. L., and D. B. Bonar. 1987. The role of DOPA and dopamine in oyster settlement behavior. *Am. Zool.* **27**: 128A.
- Coon, S. L., D. B. Bonar, and R. M. Weiner. 1985. Induction of settlement and metamorphosis of the Pacific oyster, *Crassostrea gigas* (Thunberg), by L-DOPA and catecholamines. *J. Exp. Mar. Biol. Ecol.* **94**: 211-221.
- Coon, S. L., W. K. Fitt, and D. B. Bonar. 1990. Competence and delay of metamorphosis in the Pacific oyster, *Crassostrea gigas*. *Mar. Biol.* **106**: 379-387.
- Coon, S. L., M. Walch, W. K. Fitt, D. B. Bonar, and R. M. Weiner. 1988. Induction of settlement behavior in oyster larvae by ammonia. *Am. Zool.* **28**: 70A.
- Cranfield, H. J. 1973. Observations on the behavior of the pediveliger of *Ostrea edulis* during attachment and cementing. *Mar. Biol.* **22**: 203-209.
- Crisp, D. J. 1974. Factors influencing the settlement of marine invertebrate larvae. Pp. 177-265 in *Chemoreception in Marine Organisms*, P. T. Grant, and A. M. Mackie, eds., Academic Press, London.
- Crisp, D. J., and J. S. Ryland. 1960. Influence of filming and of surface texture on the settlement of marine organisms. *Nature*. **185**: 119.
- Dube, F., and P. Guerrier. 1982. Activation of *Barnea candida* (Mollusca, Pelecypoda) oocytes by sperm or KCl, but not by  $\text{NH}_4\text{Cl}$ . requires a calcium influx. *Dev. Biol.* **92**: 408-417.
- Fitt, W. K., S. L. Coon, M. Walch, R. M. Weiner, R. R. Colwell, and D. B. Bonar. 1990. Settlement behavior and metamorphosis of oyster larvae of *Crassostrea gigas* in response to bacterial supernatants. *Mar. Biol.* **106**: 389-394.
- Gray, J. S. 1967. Substrate selection by the archianellid, *Protodrilus rubropharyngeus* Jagersten. *Helgol. Wiss. Meeresunters.* **15**: 253-269.
- Kirchman, D., S. Graham, D. Reish, and R. Mitchell. 1982. Bacteria induce settlement and metamorphosis of *Janna* (*Dexiospira*) *brasilensis* Grube (Polychaeta: Spirorbidae). *J. Exp. Mar. Biol. Ecol.* **56**: 153-163.
- Meadows, P. S., and J. I. Campbell. 1972. Habitat selection by aquatic invertebrates. *Adv. Mar. Biol.* **10**: 271-382.
- Muller, W. A. 1973. Induction of metamorphosis by bacteria and ions in the planulae of *Hydractinia echinata*: an approach to the mode of action. *Publ. Seto Mar. Biol. Lab.* **20**: 195-208.
- Neumann, R. 1979. Bacterial induction of settlement and metamorphosis in the planulae larvae of *Cassiopea andromeda* (Cnidaria: Scyphozoa, Rhizostomeae). *Mar. Ecol. Prog. Ser.* **1**: 21-28.
- Prytherch, H. F. 1934. The role of copper in the setting, metamorphosis and distribution of the American oyster, *Ostrea virginica*. *Ecol. Monogr.* **4**: 45-107.
- Roos, A., and W. F. Boron. 1981. Intracellular pH. *Physiol. Rev.* **61**: 296-434.

- Scheltema, R. S. 1961. Metamorphosis of the veliger larvae of *Nassarius obsoletus* (Gastropoda) in response to bottom sediment. *Biol. Bull.* **120**: 92-109.
- Scheltema, R. S. 1974. Biological interactions determining larval settlement of marine invertebrates. *Thalassia Jugoslav.* **10**: 263-296.
- Stevens, S. A. 1983. Ecology of intertidal oyster reefs: food, distribution and carbon/nutrient flow. Ph.D. dissertation, University of Georgia. 195 pp.
- Stoof, J. C., and J. W. Kebabian. 1984. Two dopamine receptors: biochemistry, physiology and pharmacology. *Life Sci.* **35**: 2281-2296.
- Thorson, G. 1950. Reproductive and larval ecology of marine bottom invertebrates. *Biol. Rev.* **25**: 1-45.
- Ward, S., E. Hogan, and G. A. Nelson. 1983. The initiation of spermiogenesis in the nematode *Caenorhabditis elegans*. *Dev. Biol.* **98**: 70-79.
- Weiner, R. M., A. M. Segall, and R. R. Colwell. 1985. Characterization of a marine bacterium associated with *Crassostrea virginica* (the Eastern oyster). *Appl. Environ. Microbiol.* **49**: 83-90.
- Weiner, R. M., M. Walch, M. P. Labare, D. B. Bonar, and R. R. Colwell. 1989. Effects of biofilms of the marine bacterium *Alteromonas colwelliana* (LST) on set of the oysters *Crassostrea gigas* (Thunberg, 1793) and *C. virginica* (Gmelin, 1791). *J. Shellfish Res.* **8**(1): 117-123.
- Williams, G. B., E. M. Elder, and M. Sussman. 1984. Modulation of the cAMP relay in *Dictyostelium discoideum* by ammonia and other metabolites: possible morphogenetic consequences. *Dev. Biol.* **105**: 377-388.
- Wilson, D. P. 1955. The role of micro-organisms in the settlement of *Ophelia bicornis* Savigny. *J. Mar. Biol. Assoc. U. K.* **34**: 531-543.
- Zar, J. H. 1974. *Biostatistical Analysis*. Prentice-Hall, New Jersey. 620 pp.
- Zobell, C. E., and E. C. Allen. 1935. The significance of marine bacteria in the fouling of submerged surfaces. *J. Bacteriol.* **29**: 239-251.